



CONTENTS

[Highlights](#)

[What is vinyl chloride?](#)

[What happens to vinyl chloride when it enters the environment?](#)

[How might I be exposed to vinyl chloride?](#)

[How can vinyl chloride affect my health?](#)

[How likely is vinyl chloride to cause cancer?](#)

[How can vinyl chloride affect children?](#)

[How can families reduce the risk of exposure to vinyl chloride?](#)

[Is there a medical test to show whether I've been exposed to vinyl chloride?](#)

[Has the federal government made recommendations to protect human health?](#)

[References](#)

[Contact Information](#)

RELATED RESOURCES

[ToxFAQ™](#)  29k

[Public Health Statement](#)  46k

[Toxicological Profile](#)  1.9MB

[MMG](#)  54k

September 2004

ToxFAQs™ for Vinyl Chloride

CAS# 75-01-4

This fact sheet answers the most frequently asked health questions about vinyl chloride. For more information, you may call the ATSDR Information Center at 1-888-422-8737. This fact sheet is one in a series of summaries about hazardous substances and their health effects. This information is important because this substance may harm you. The effects of exposure to any hazardous substance depend on the dose, the duration, how you are exposed, personal traits and habits, and whether other chemicals are present.

HIGHLIGHTS: Exposure to vinyl chloride occurs mainly in the workplace. Breathing high levels of vinyl chloride for short periods of time can cause dizziness, sleepiness, unconsciousness, and at extremely high levels can cause death. Breathing vinyl chloride for long periods of time can result in permanent liver damage, immune reactions, nerve damage, and liver cancer. This substance has been found in at least 616 of the 1,647 National Priority List sites identified by the Environmental Protection Agency (EPA).

What is vinyl chloride?

Vinyl chloride is a colorless gas. It burns easily and it is not stable at high temperatures. It has a mild, sweet odor. It is a manufactured substance that does not occur naturally. It can be formed when other substances such as trichloroethane, trichloroethylene, and tetrachloroethylene are broken down. Vinyl chloride is used to make polyvinyl chloride (PVC). PVC is used to make a variety of plastic products, including pipes, wire and cable coatings, and packaging materials.

Vinyl chloride is also known as chloroethene, chloroethylene, and ethylene monochloride.

[back to top](#)

What happens to vinyl chloride when it enters the environment?

- Liquid vinyl chloride evaporates easily. Vinyl chloride in water or soil evaporates rapidly if it is near the surface.

[an error occurred while processing this directive]

ATSDR RESOURCES

[ToxFAQs™](#)

[ToxFAQs™ en Español](#)

[Public Health Statements](#)

[Toxicological Profiles](#)

[Minimum Risk Levels](#)

[MMGs](#)

[MHMIs](#)

[Interaction Profiles](#)

[Priority List of](#)

[Hazardous Substances](#)

[Division of Toxicology](#)

- Vinyl chloride in the air breaks down in a few days to other substances, some of which can be harmful.
- Small amounts of vinyl chloride can dissolve in water.
- Vinyl chloride is unlikely to build up in plants or animals that you might eat.

[back to top](#)

How might I be exposed to vinyl chloride?

- Breathing vinyl chloride that has been released from plastics industries, hazardous waste sites, and landfills.
- Breathing vinyl chloride in air or during contact with your skin or eyes in the workplace.
- Drinking water from contaminated wells.

[back to top](#)

How can vinyl chloride affect my health?

Breathing high levels of vinyl chloride can cause you to feel dizzy or sleepy. Breathing very high levels can cause you to pass out, and breathing extremely high levels can cause death.

Some people who have breathed vinyl chloride for several years have changes in the structure of their livers. People are more likely to develop these changes if they breathe high levels of vinyl chloride. Some people who work with vinyl chloride have nerve damage and develop immune reactions. The lowest levels that produce liver changes, nerve damage, and immune reaction in people are not known. Some workers exposed to very high levels of vinyl chloride have problems with the blood flow in their hands. Their fingers turn white and hurt when they go into the cold.

The effects of drinking high levels of vinyl chloride are unknown. If you spill vinyl chloride on your skin, it will cause numbness, redness, and blisters.

Animal studies have shown that long-term exposure to vinyl chloride can damage the sperm and testes.

[back to top](#)

How likely is vinyl chloride to cause cancer?

The U.S. Department of Health and Human Services has determined that vinyl chloride is a known carcinogen. Studies in workers who have breathed vinyl chloride over many years showed an increased risk of liver cancer; brain cancer, lung cancer, and some cancer of the blood have also been observed in workers.

[back to top](#)

How can vinyl chloride affect children?

It has not been proven that vinyl chloride causes birth defects in humans, but studies in animals suggest that vinyl chloride might affect growth and development. Animal studies also suggest that infants and young children might be more susceptible than adults to vinyl chloride-induced cancer.

[back to top](#)

How can families reduce the risk of exposure to vinyl chloride?

Tobacco smoke contains low levels of vinyl chloride, so limiting your family's exposure to cigarette or cigar smoke may help reduce their exposure to vinyl chloride.

[back to top](#)

Is there a medical test to show whether I've been exposed to vinyl chloride?

The results of several tests can sometimes show if you have been exposed to vinyl chloride. Vinyl chloride can be measured in your breath, but the test must be done shortly after exposure. This is not helpful for measuring very low levels of vinyl chloride. The amount of the major breakdown product of vinyl chloride, thiodiglycolic acid, in the urine may give some information about exposure. However, this test must be done shortly after exposure and does not reliably indicate the level of exposure.

Vinyl chloride can bind to genetic material in your body. The amount of this binding can be measured by sampling your blood and other tissues. This measurement will give information about whether you have been exposed to vinyl chloride, but it is not sensitive enough to determine the effects on the genetic material resulting from exposure. These tests are not available at most doctors' offices, but can be done at special laboratories.

[back to top](#)

Has the federal government made recommendations to protect human health?

Vinyl chloride is regulated in drinking water, food, and air. The EPA requires that the amount of vinyl chloride in drinking water not exceed 0.002 milligrams per liter (mg/L) of water.

The Occupational Safety and Health Administration (OSHA) has set a limit of 1 part vinyl chloride per 1 million parts of air (1 ppm) in the workplace.

The Food and Drug Administration (FDA) regulates the vinyl chloride content of various plastics. These include plastics that carry liquids and plastics that contact food. The limits for vinyl chloride content vary depending on the nature of the plastic and its use.

[back to top](#)

References

Agency for Toxic Substances and Disease Registry (ATSDR). 1997. Managing Hazardous Materials Incidents. Volume III – Medical Management Guidelines for Acute Chemical Exposures: [Vinyl Chloride](#). Atlanta, GA: U.S. Department of Health and Human Services, Public Health Service.

Agency for Toxic Substances and Disease Registry (ATSDR). 2004. [Toxicological Profile for Vinyl Chloride](#) (*Draft for Public*

Comment). Atlanta, GA: U.S. Department of Health and Human Services, Public Health Service.

[back to top](#)

Where can I get more information?

ATSDR can tell you where to find occupational and environmental health clinics. Their specialists can recognize, evaluate, and treat illnesses resulting from exposure to hazardous substances. You can also contact your community or state health or environmental quality department if you have any more questions or concerns.

For more information, contact:

Agency for Toxic Substances and Disease Registry
Division of Toxicology
1600 Clifton Road NE, Mailstop F-32
Atlanta, GA 30333
Phone: 1-888-42-ATSDR (1-888-422-8737)
FAX: (770)-488-4178
Email: ATSDRIC@cdc.gov

[back to top](#)

ATSDR Information Center / ATSDRIC@cdc.gov / 1-888-422-8737

This page was updated on October 15, 2004

[ATSDR Home](#) | [Search](#) | [Index](#) | [Glossary](#) | [Contact Us](#)
[About ATSDR](#) | [News Archive](#) | [ToxFAQs](#) | [HazDat](#) | [Public Health Assessments](#)
[Privacy Policy](#) | [External Links Disclaimer](#) | [Accessibility](#)
[U.S. Department of Health and Human Services](#)